

[SQUEAKING]

[RUSTLING]

[CLICKING]

CASEY So I'm going to prove a few theorems about limits, which will allow us to compute limits, or at least we can use to
RODRIGUES: prove that other non-trivial limits exist using these theorems, rather than using the definition directly.

So this first theorem is the easiest theorem in the world because it's simply just restating the definition of convergence of a sequence. So I'm going to state it as follows, so pretty short, that if I have a sequence x_n , then it converges to x if and only if the sequence obtained by taking the absolute value of $x_n - x$ equals 0, or the limit of that sequence is 0.

So what is the proof? It follows just immediately from the definition. So I'm not even going to write anything. I'll leave it to you. But the proof follows from the definition and the simple fact that $|x_n - x|$ in absolute value is equal to the absolute value of the absolute value of $x_n - x$ minus 0, OK.

So the definition says for all ϵ positive, you have to find an M so that this is less than ϵ for all n bigger than or equal to capital M . But if you found such a capital M for this to be less than ϵ , then this will be less than ϵ , which is saying that this limit equals 0.

And then going the other direction, it's the same thing. So this is just following directly from the definition and this simple fact. OK, that was a very silly fact about limits, but a very useful one in conjunction with the next theorem, which is not so trivial, which is the squeeze theorem.

And it says the following. So let a_n , b_n , and x_n be sequences such that the following holds for all n natural numbers, $a_n \leq x_n \leq b_n$. And a_n and b_n are convergent sequences.

And their limits equal each other. And they're given by some number, call it x . Then the conclusion is that the limit as n goes to infinity of x_n equals x . So when I write something like this, you should also kind of-- there's half a sentence before that that goes with this saying, x_n is a convergent sequence. And its limit is equal to x . So it's two statements in one when I write that.

So if you just draw a picture, the squeezed theorem shouldn't be too surprising. So this is a little discussion. So here's x , the common limit of a_n and b_n . And so we can imagine that we're trying to show the limit as n goes to infinity of x_n equals x .

So that means we have to find for every ϵ a capital M so that x_n is between $x + \epsilon$ and $x - \epsilon$. So if I go out a little bit, I would hope I can find a natural number, a capital M so that $x_n - x$ or x_n is in this little interval.

Now, if I'm assuming that a_n and x_n , if a_n and b_n are squeezing x_n , in other words x_n is between the two, and b_n is converging to x , then for n bigger than or equal to some integer M_0 , all of the b_n 's are in this interval. OK, maybe they're not there. Maybe they could be over here. But the way I drew it is just to the right of x .

And, likewise, since a_n is converging to x , there exists some other integer M_1 so that if I look at a_n , it's also in this interval. Maybe it's over here. Maybe it's-- well, it can't be to the right of b_n because that inequality up there strictly implies that a_n is less than or equal to b_n . But it's in this interval.

So then what would that say if I look at n bigger than or equal to $n_0 + M_1$, $M_0 + M_1$, then n is bigger than M_0 , this guy. And n is bigger than both and M_1 , this guy. And therefore if I look at x_n , it's going to be between these two. And in particular, it's going to be in this interval.

So that's the proof in a picture. And now our goal is just to write it down. So that's the picture of the proof. But if you're actually trying to guess why this would be true? I mean, the b_n 's, you can imagine, are getting very close to x . The a_n 's are also getting very close to x . x_n is in between the two, so it's getting squeezed to x , and thus the name.

OK, so now we just need to turn this into written word. So we need to show-- we're going to show that x_n converges to x . So we're-- all we have is an epsilon delta proof. Or we could use that theorem. But let's go with an epsilon delta proof. I mean, not epsilon delta, epsilon M . Epsilon delta proofs will come later.

So let epsilon be positive. And since b_n converges to x , there exist to M_0 in natural numbers such that for all n bigger than or equal to n_0 , $b_n - x$ is less than epsilon in absolute value, which is the same as saying-- well, I mean it's not the same. But it implies that b_n is less than $x + \epsilon$.

Since a_n 's converge to x as well, there exists a M_1 , natural number, such that for all n bigger than or equal to M_1 , $a_n - x$ in absolute value is less than epsilon, which implies a_n is between $x - \epsilon$ and $x + \epsilon$, but I'm only going to use one of those inequalities.

Now I'm going to choose the capital M for my sequence x_n , I'm trying to show convergence to x . So choose m to be $n_0 + M_1$. I mean, you could have chosen it to be the maximum of the two. But this works just fine as well.

Then if n is bigger than or equal to m , this implies n is bigger than or equal to M_0 and M is bigger than or equal to M_1 , which implies that both of these inequalities here are valid for this n . So then $x - \epsilon$ is less than a_n , which by assumption, is less than or equal to x_n , which is less than or equal to b_n , which is less than $x + \epsilon$.

Now, these string of inequalities, therefore, tell us that $x - \epsilon$ is less than x_n is less than $x + \epsilon$, which is equivalent to saying the absolute value of $x_n - x$ is less than epsilon. And therefore x_n converges to x .

So these two facts together give us a very robust and short way to prove limits of sequences. So, for example, let me give you a simple one. Let's show the limit as n goes to infinity of n^2 over $n^2 + n + 1$ equals 1. So, I mean, this is a very simple limit to use these theorems on. But in practice, you don't always have just a simple expression like this.

And we'll use these two theorems in conjunction to prove some other theorems here in a minute. But let's just see it in action once. Let me look at-- so this sequence converges to 1 if and only if the absolute value of the difference converges to 0. So let's look at the absolute value of the difference. This is equal to-- now just doing the algebra-- this is equal to-- and taking absolute values gives me $n + 1$ over $n^2 + n + 1$.

And this is less than or equal to. This 1 is making things bigger, so I can drop it. And this is less than or equal to $n + 1$ over $n^2 + n$. Now, $n^2 + n$ I can factor into n times $n + 1$. So then that cancels with this $n + 1$ on top. And I just get $1/n$.

So 0 is less than or equal to $n^2 / (n^2 + n + 1) - 1$, which is less than or equal to $1/n$. Now, $1/n$ we've shown using epsilon delta proof-- I mean, epsilon M proof. We've shown $1/n$ converges to 0.

So since 0 converges to 0, the left side, and $1/n$ converges to 0, the right side, that implies by the squeeze theorem, $n^2 / (n^2 + n + 1) - 1$ converges to 0 by the squeeze theorem. Which implies that $n^2 / (n^2 + n + 1)$ converges to 1 by that first theorem.

OK now you can imagine that instead of having this at your disposal, I asked you to do an epsilon proof, an epsilon M proof of this statement, then you would have taken this and played with it just like we did here and gotten to $1/n$. And therefore if this is less than epsilon, that would imply this is less than epsilon. So you would choose capital M to be-- so that $1/\text{capital M}$ is less than epsilon. But using these two theorems saves us a little work and a little time.

OK, now, so at the end of the lecture last time, we discussed the notion of subsequences. And we showed that limiting-- limits and subsequences interact nicely, meaning if I have a convergent sequence, then every subsequence converges to the same thing.

So now a natural question is, how do limits interact with the order of the real numbers? $\mathbb{R}$ has these two fundamental properties about it, that it's-- so first off, that it has the least upper bound property, and also that it's an ordered field.

So first natural question is, how does this definition of limit convergence interact with the order? So the first theorem states, or this term that kind of answers this question is the following is that limits respect order, basically. So if x_n, y_n , are convergent sequences and for all n , x_n is less than or equal to y_n , then what should be the conclusion?

The limit as n goes to infinity. So what I said a minute ago as limits respect inequality. Then I should be able to take the limits of both sides and still have this inequality. Then limit as n goes to infinity is less than or equal to limit as n goes to infinity of y_n .

And a simple corollary that follows from this is that if x_n is a convergent sequence, and for all n a natural number, you have two numbers a and b , such that a_n is less than or equal to-- or a is less than or equal to x_n is less than or equal to b , then this implies that the limit of x_n is also between a and b .

Let me say something very brief about what this says and doesn't say. So what does this not say? You may lose a strict inequality, meaning what? It can be the case that x_n is less than y_n for all of n , but the limit of x_n equals the limit of y_n .

So simply having less than x_n less than y_n does not imply the limit of x_n is less than the limit of y_n . All right, so what do I mean by this? This does not imply that the limit is less than y_n .

Now, at this point in class, I would ask somebody to give me a counter example. So at this point, I'm going to take a bite of my cookie and let you think about that. I didn't have to take a bite of my cookie. You could have just paused the video and thought about it. But then I wouldn't get a bite of my cookie.

OK, so what's an example of two sequences that satisfy this, but don't satisfy this? If x_n equals 0 for all n , and y_n equals $1/n$, then the x_n is less than y_n for all n . And what's the limit x_n ? Well, that's just 0.

And before the limit of y_n , that's just 0. And these two things equal each other. This is not less than that, OK. So just want to make that small point. Could pop up as a small question on one of the midterms-- I should say the midterm-- and possibly the final.

All right, so let's prove-- [COUGHS] let's prove 1. 2 follows immediately from one. For two, we simply take, for example, the upper inequality x_n is less than or equal to b , we take y_n to be the constant sequence b .

And for the other one, we would take the bigger sequence to be x_n , and the smaller sequence to be the constant sequence a . So two follows immediately from 1, so we're just going to do 1. And we haven't done a proof by contradiction in a while, so why not do it by contradiction?

OK, so the proof-- let's label these sequences so I don't have to write limit as n goes to infinity, and limit of x_n , and limit as n goes to infinity of y_n so much. So let's call their limits x and y . And what do we want to show? x is less than or equal to y .

That's our goal. And we're going to prove this by contradiction. That is, let's assume y is less than x , and arrive at a false statement, which contradicts our setting our assumptions that we have. Assume y is less than x . Now, let me draw a picture here to go along with what's going to happen.

So if y is less than x , all of the y_n 's have to be near y if I go far enough out. And all the x_n 's have to be near x as long as I go far enough out. So that would contradict eventually the fact that the x_n 's are supposed to be less than or equal to the y_n 's, if y is less than x .

And let's say I go out, let's say, half the distance between y and x . So this is like the midpoint. And all the y_n 's are here, and all the x_n 's are here, and the y_n 's are here, then I cannot have x_n less than or equal to y_n , which is my assumption, which is the assumption in my theorem.

Now, maybe I should have written this here. Sometimes it's good to reiterate what your assumptions are. Suppose for all n , x_n is less than or-- y_n . And-- like that. And so the thing we're trying to show is this, OK. So we assume the negation of this and arrive at a contradiction to the things we're assuming, or just to general true facts.

I had to add the word true on the facts, because they're, like I said, at some point there's alternative facts flying around out there. OK, so this picture, we're going to turn this into a proof. So since y_n converges to y , there exists a natural number $M \geq 0$ such that for all n bigger than or equal to M , $y_n - y$ is less than $x - y$.

Now, that's a positive number, because we're assuming x is bigger than y . And by the definition of limit, given any positive number, I can find an integer so that for all n bigger than or equal to that integer, this thing is less than that small number. And I'm just choosing that small number to be this very special small number, because that's going to help me arrive at a contradiction.

Since x_n converges to x , there exist M_1 natural number such that for all n bigger than or equal to M_1 , $x_n - x$ is less than the same thing. OK, so this is putting into a precise form what I was saying that all the x_n 's to be close to x eventually. And all the y_n 's to be close to y eventually. And how eventually? Well, eventually enough so that I'm in these two disjoint intervals, OK.

Let n be $M_0 + M_1$. Then n is bigger than or equal to M_0 . And n is bigger than or equal to M_1 . So both of these inequalities are valid for this n and what does this mean? Well, then that implies that $y_n - x_n$ so let me tack on one more inequality.

Just by removing the absolute values and adding y , this tells me y_n is less than $x + y/2$. And this tells me x_n is less than-- or the other way, sorry. Well, we'll just write this down here. So

Then y_n is less than $y + x - y/2$, which equals $x + y/2$, which equals $x - x_n + x_n + y/2$. And this is less than x_n by the second inequality. So this follows from the first inequality. This follows from the second inequality, which implies for this specific n , y_n is less than x_n .

And this is a contradiction to our assumption that y_n is bigger than or equal to x_n for all n . So we had just found, based on this assumption here, that we arrive at a contradiction to our other assumptions. And therefore this must be false.

So that's how limits interact with inequalities. So if I have two sequences, one bigger than the other, then the limits respect that inequality. That has to deal with the order part of $\mathbb{R}$ being an ordered field. So what about the field part of $\mathbb{R}$ being an ordered field? So how does limits interact with algebraic operations?

All right, quite well, it turns out. So let's in theorem. So suppose I have two convergence sequences, limit as n goes to infinity of $x_n = x$, and limit as n goes to infinity of $y_n = y$, then several things hold.

The first is that, again, you should kind of read this as two statements written in one, limit as n goes to infinity of $x_n + y_n$. So this is a new sequence that I formed by just taking the term-by-term sum of these two things. This sequence, this new sequence is convergent. And the limit equals the sum of the limits, OK?

The second is, for all c in $\mathbb{R}$, the limit as n goes to infinity of the new sequence obtained by taking every entry of the sequence x_n and multiplying it by this fixed number c , the limit of that product is the product of c and x . So limits respect what one would call scalar multiplication.

But we could be more general than that. c you can think of as just one example of a convergent sequence, just a constant sequence. But, in general, we have that the product of two convergent sequences is convergent. And the limit of the product is the product of the limit.

And, finally, if we have something for-- write it over here-- if we have something for a product, then perhaps we have something by quotient. And that's as long as we can divide things. So if for all n , y_n does not equal 0, and the limit y does not equal 0, then limit of the quotient x_n / y_n is the quotient of the limits, OK.

OK, so we're going to prove this first one using this scheme of using both this simple theorem about limits and the squeeze theorem. And it's quite simple. So we just use the triangle inequality. By the triangle inequality, 0 is less than or equal to $x_n + y_n - x - y$. And $x_n - x$, $y_n - y$.

And then I use the triangle inequality. I get $x_n - x + y_n - y$ -- oh -- you know what, scratch that. Because, in fact, I was trying to be too clever and would have ended up using the theorem to prove the theorem. So let's not do that. It's never good to use the theorem you're trying to prove to prove the theorem that you're trying to prove.

So let's go back to basics and use the definition. Let ϵ be positive. So since x_n converges to x , there exists a natural number M such that for all n bigger than or equal to M , $x_n - x$ -- it's supposed to be a n , but it looks like a k -- less than $\epsilon/2$. Why the 2? Well, you'll see in a minute.

And similarly for the sequence y , there exist M_1 natural number such that for all n bigger than or equal to M_1 , $y_n - y$ is less than $\epsilon/2$. So I have these two integers, which are given to me by the fact that x_n converges to x , y_n converges to y , and the definition of convergence.

I can always find these two integers for any small tolerance. And I'm choosing the tolerance to be $\epsilon/2$ for some reason, which you'll see in a minute. And so what is the integer capital M that I choose for this ϵ for the sequence $x_n + y_n$?

I'm going to choose M to be $M_0 + M_1$. And now I need to show that this choice of M works. And if n is bigger than or equal to M , this implies that n is bigger than or equal to M_0 . And M is bigger than or equal to M_1 . So both of these inequalities are valid for this n . And therefore I get $x_n + y_n - x - y$.

And now I do what I was going to do a minute ago when I was going to use the theorem to prove a theorem. $x_n - x$, $y_n - y$, I group those together. And then I used a triangle inequality. This is less than or equal to $x_n - x + y_n - y$.

And now this is less than $\epsilon/2$. This is less than $\epsilon/2$, so equals ϵ . And now you can see why I chose the 2. Because I wanted to show this was less than ϵ . And I had control over these two things. And the sum of controls gives me ϵ . So I choose the control to be $\epsilon/2$.

If I had three sequences, then you could probably guess though which is $\epsilon/3$. Meaning if I had sequences x_n , y_n , and z_n , and I looked at $x_n + y_n + z_n$, I could show that converges to the sum of the limits. And I would choose these integers M_0 , M_1 , M_2 , so that I have $\epsilon/3$ here, so that they sum up to ϵ .

So now we prove 2, that for this single scalar multiplication if you like where you just multiply each term by a single number, the limit respects that multiplication. Do an ϵ proof again. Since x_n converges to x -- so now we're trying to show that second limit -- there exists M_0 , a natural number, such that for all n than or equal to M_0 , $x_n - x$ is less than $\epsilon/|c| + 1$, OK.

And now you have to trust me, why that thing? Well, you'll see. It'll come out just like this did. So for the sequence $c \cdot x_n$, we'll choose M to be just this M_0 . Then if n is bigger than or equal to M , which is equal to M_0 , this implies this inequality holds. And therefore $c \cdot x_n - c \cdot x$.

This c pops out of the absolute value and becomes the absolute value of c times the absolute value of $x_n - x$. And which is less than-- so I'm writing less than here. But this thing is less than that. [INAUDIBLE] is less than c over $c + 1$ epsilon.

Now, this quotient here, this number over this number plus 1, is always less than 1. So this positive number which is less than 1 or non-negative number which is less than 1 is going to be less than 1 times epsilon, which gives me epsilon.

And maybe you're wondering why didn't I choose-- so this is just a little smidgen of sophistication, not much, just a little. Why didn't I choose this so that it's epsilon over the absolute value of c so that when I stick in my inequality for this guy I just get epsilon?

Well, what if c equals 0? So then I'm telling you to choose capital M_0 so that the absolute value of $x_n - x$ is less than epsilon over 0. Division by 0 is a no no. But if we fudge it a little by adding 1 we get something that still does the job. It still gives me some number which is non-negative and less than 1, which is enough, OK.

So let's prove that the limit of the product is the product of the limits. Since the sequence y_n converges to y , it's a convergent sequence, and therefore it's bounded. That means that there exists some non-negative real number b such that for all natural numbers n , y_n is less than or equal to b in absolute value.

Then I look at $x_n y_n - x y$ and I add and subtract $x y_n$. You can write this as plus. And now I use the triangle inequality that this is less than or equal to $|x_n - x| y_n$ plus $|y_n - y| x$. And y_n is bounded by b for all n . So this is less than or equal to-- plus $|x_n - x| b$ plus $|y_n - y| x$.

Now, let me just state the obvious that we get 0 is less than or equal to $|x_n - x| b$ plus $|y_n - y| x$. And this is less than or equal to, as we've shown here, plus $|x_n - x| b$ plus $|y_n - y| x$. Now, the right-hand side-- so the left side of this inequality converges to 0. The right side also converges to 0 because we've just shown by 1 and 2.

By 2, this converges to 0. And by that theorem, since x_n converges to x , this product here converges to 0. b is a fixed number. And the same thing for this, that converges to 0. And therefore by 2, this sum converges to 0. OK, so these two arrows are by 2. And this is by 1.

So let me just summarize by 1 and 2 the right-hand side of this inequality, $b|x_n - x|$ plus $|y_n - y|x$. This converges to 0. Which by the squeeze theorem implies that-- this is by squeeze theorem-- which implies that $x_n y_n$ converges to $x y$ by that first theorem.

So I'm not going to keep referring to that first theorem. Because it's such a simple fact, I'm just going to keep using it without referencing it, namely that the sequence, a sequence converges to x if and only if the absolute value of this thing, this difference, converges to 0, OK.

All right, so that proves the limit of the product is the product of the limits. Now, for the quotient we can use 3 once we've proven it for just 1 over y_n . So what do I mean? So now we're assuming the y_n is not equal to 0 for all n . And y does not equal 0.

So if we prove this statement that the limit as n goes to infinity of $1/y_n$ equals $1/y$, then by 3 implies that the limit as n goes to infinity of x_n/y_n equals x/y . Because x_n/y_n is just a product of x_n and $1/y_n$. So we just need to prove this special case if you like.

And we're to do it kind of the same way. Now because we're dividing by y_n , so here we use that we had an upper bound for the product here. But when we take $1/y_n$ to get an upper bound of that, it means we need a lower bound on y_n , on the absolute values of y_n . And we get that by our assumption that the limit is non-zero for all n , and so is the y_n 's.

So first thing we prove, or let me write this as a claim, there exists a positive number b , little b , such that for all natural numbers n , y_n is bigger than or equal to b . And we know that a sequence is bounded. So we know that there's always a capital B so that absolute value of y_n is less than or equal to B .

But for sequences that are non-zero and converging to a non-zero limit, then you can bound them away from 0. And it's kind of the same proof that we gave for showing that a convergent sequence is bounded above. So let me draw a quick picture. And let's assume that the limit is positive just for the sake of the picture.

So this is a little discussion on why this is true. And this picture is going to look-- at least the explanation is going to be kind of similar to why a sequence is bounded. And this one is why is it bounded below. So let's assume that the limit y is positive.

And let's say I go out within distance, let's say, $y/2$ so that I'm still positive. So this is y minus $y/2$, absolute value. So in this picture y is positive. So that's just equal to $y/2$. Then what can I say? That eventually all of the y_n 's are here in this interval away from 0. And, in fact, their absolute value is bounded by $y/2$. So let me just write it that way.

So all the y_n 's or n bigger than or equal to some M , all have to lie in this interval because they're converging to y and y is positive. And therefore in absolute value, they're all bounded above-- below I mean-- by $y/2$. They're all at least distance $y/2$ to 0.

OK, and then all that's left to handle are-- maybe the finitely many that are left, y_1, y_2, y_m minus 1, that are scattered on the real line but are non-zero. So we're just going to end up taking the minimum of this number and the absolute value of these numbers.

So since the y_n 's converge to y , and y does not equal 0, there exist an integer M such that all n bigger than or equal to capital M . So this picture was why this claim is true. It was not the proof of why this claim is true. What I'm writing now is the actual proof of why this claim is true-- such that for all n bigger than or equal to a capital M , y_n minus y in absolute value is less than $y/2$.

So for this picture I drew where y is positive. That would have just been $y/2$. But you have to use an absolute value for the other case that y is negative, because this has to be a positive number. Then for all n bigger than or equal to M , this any inequality and the triangle inequality gives me-- so if I look at the absolute value of y , this is equal to the absolute value of y minus y_n plus y_n .

And now I use a triangle inequality, which is less than-- this is less than $y/2$ plus y_n . And I started off with the absolute value of y . So when I subtract that over, that tells me that absolute value of $y/2$ is less than absolute value of y_n for all n bigger than or equal to capital M .

So then I let b to be the minimum of several numbers. Now I'm writing $\min$, but I should write $\inf$. But so if you like, let me write $\inf$ of $y_1, y_m - 1$, and y over 2 . And by what you are doing on the assignment, I think it was the assignment 2, this $\inf$ always exists in a finite set.

This is a finite set of positive numbers. And therefore its infimum exists as one of these elements. One of these M numbers is the infimum. And they're all positive, so this is a positive number. And so then simply how this number is defined, it follows that for all n , y sub n is bigger than or equal to b .

Because, again, if n is between-- little n is between 1 and capital $M - 1$, then certainly that the absolute value of that thing is bigger than or equal to the smaller of all of these, which is bigger than or equal to b . And if n is bigger than or equal to capital M , then we proved over here that y sub n is bigger than the absolute value of y over 2 , which is bigger than or equal to the minimum of these numbers and y over 2 , which equals b .

All right, so that proves the claim. That proves the claim. But we haven't proved what we wanted to yet that the limit as n goes to infinity of 1 over y sub n equals 1 over y . But this follows almost immediately from what we've done so far.

So now we're going to use the claim to prove it. So we look at-- compute that the 0 is less than or equal to 1 over y sub n minus 1 over y . We're going to show that this goes to 0 using those two theorems. And so by algebra that's equal to-- and using the absolute value, this is 1 over y sub n minus y . I mean, the absolute value of y sub n minus y over absolute value of y sub n absolute value of y .

Now y sub n is bigger than or equal to b . So this is less than or equal to 1 over b times y times y sub n minus y . So just to summarize, we've shown that 0 is less than 1 over y sub n minus 1 over y is less than 1 over b times 1 over-- b times the absolute value of y times y sub n minus y .

Now, this goes to 0 because it's just a constant sequence. This converges to 0 because y sub n minus y in absolute value goes to 0 . This is just a fixed number of times that. And by what we've proven for 2, this product converges to 0 .

So by the squeeze theorem, we get that 1 over y sub n , this converges to 0 , which implies-- OK. So another big property about the real numbers that we proved after we stated the existence of the real numbers, which remember was this is defined as this ordered field with a least upper bound property.

We proved that the square root of 2 exists as a real number. There was really nothing special about 2 . In fact, you could prove that the square root of x exists as a real number for any x that's a positive or non-negative number.

So the square root of a real number is well defined. Or the square root of a non-negative number is well defined and always exists as a real number. So you can ask how does limits interact with square roots? And they interact just as you think they should.

If I have a sequence so that for all n , x_n is bigger than or equal to 0, and it's a convergent sequence, converging to some number x , then the limit of the square roots of these guys equals the square root of the limit, OK.

Now, I want you to just take a second here and understand that this is a meaningful statement. Because since the x_n 's are all non-negative by a theorem that-- let's see, did I erase it already? The one that had to deal with limits and the order, so since the x_n 's are all non-negative, that implies that x is non-negative so that the square root is meaningful.

OK, so first check whenever somebody says here's this theorem, or this theorem is-- I think this theorem is true-- is to check to make sure that a theorem is meaningful. So let's prove this. So there's two cases to consider, x is equal to 0 or x is non-zero. So let's do the first case.

So the limit is 0. So we'll do this proof using the definition of limits, meaning the epsilon M definition. So we want to show that the limit of the square root of x_n equals 0. So let epsilon be positive. And since x_n converges to 0, there exists natural number n_0 such that if n is bigger than or equal to n_0 , then x_n minus the limit, which is just 0, and taking the absolute value, which is just x_n , which is equal to x_n because x_n is non-negative, is less than epsilon squared.

Remember, I can always find, no matter what is underneath my hand, since x_n converges to 0, I can find a natural number so that that thing is less than what's underneath my hand. And the thing that I'm going to have underneath my hand that's going to make things work out for the square root is epsilon squared, OK.

Choose M to be n_0 . So I'm going to show this M works for the sequence square root of x_n . And n bigger than or equal to M . The square root of x_n minus 0, which is just x_n .

Now, it's also-- so we didn't strictly speaking prove this-- but it's not too hard to show that square roots respect inequalities. So x_n is less than epsilon squared. So the square root is less than epsilon squared, which equals epsilon.

So the second case is x not equal to 0. And to do this case, we'll use those two theorems again. And let's look at square root of x_n minus square root of x . Now, if I write this as-- and multiply top and bottom by x_n plus square root of x , square root of x_n plus square root of x , which is a positive number. It's fine to divide by it as well, because x is non-zero.

So here not just non-zero, it's positive. Because x has to be non-negative. Now this is the product of something minus something else with something plus something else. So then that's going to be the difference of the squares. So that's equal to-- over-- and, again, these are positive numbers so that they come out of the absolute value.

And this is non-negative. So, in fact, it's only making things bigger on the bottom and therefore things smaller overall. So this is less than or equal to-- just if I replace this by square root of x . So what did I prove? 1 over square root of x -- all right.

And so by assumption, x_n converges to x . So this goes to 0. This is a fixed number multiplied by this thing going to 0. So by number 2 and the theorem we proved before, this whole product converges to 0. So, and of course this converges to 0. So this thing in the middle must converge to 0 by the squeeze theorem.

So that's the square root. And let me just remark that-- so this number 3 up here, that the limit of the product is a product of the limits, this thus implies that the limit of x_n squared converges to x squared. So the square of x_n converges to the square of the limit. And by induction, you can show that the cube fourth power, fifth power of x_n converges to the fourth power, fifth power of x .

And not only that, you can also prove-- so I'm not going to do this. And I'm not going to force you to either. But just know these as facts, that I don't have to just take the square root, I could take the k -th root. And this statement still is true that if x_n is bigger than or equal to 0 for all n , and I have a limit, then the k -th root of x_n converges to the k -th root of x .

So the final theorem we'll prove for today, which will conclude our facts about limits, is we have-- I mean, we've been using this all along, although I haven't made special attention about it. The real numbers there, this, again, like I said, an ordered field with a least upper bound property.

So we've seen how limits interact with this structure of the real numbers. But they also have a distance associated to them, the absolute value. The distance from two numbers a and b is the absolute value of a minus b . Or the distance from a number to 0 is just the absolute value of that number.

So one could ask, how does the limit interact with this additional structure of the absolute value? And so just like everything's been good so far with limits, it's the, same with the absolute value namely that limits respect absolute value. So if x_n is convergent sequence, the limit x , then the sequence of absolute values is also a convergent sequence.

And the limit as n goes to infinity of the absolute values equals the absolute value of the limit. Now let me, again, let's try to think a little more deeply about this real quick. If I have a convergence sequence and the absolute values converge, does the converse hold?

If the absolute values converge, does this imply that the original sequence converges? And the answer is, of course, no. Well, not of course. I didn't even give you a minute to think about it. But why is the converse not true? So let me make this a remark.

Converse is not true because you could look at x_n equals minus 1 to the n . Then the absolute values of these guys converges, but the original sequence does not converge. So this is a one-way street for convergence and the convergence of the absolute values.

So before I prove this theorem, let me just prove a quick inequality which I think I said I was going to put on the assignment, and then forgot to put on the assignment. But you should know it. So this theorem is the reverse triangle inequality which states for all a, b real number, the absolute value of the difference in absolute values is less than or equal to the absolute value of the difference.

OK, so the proof of this reverse triangle inequality just uses the original triangle inequality. So absolute value of a , this is equal to the absolute value of a minus b plus b . And this is less than or equal to the absolute value of a minus b plus absolute value of b . And thus the absolute value of a minus the absolute value of b is less than or equal to the absolute value of a minus b .

Now, these are just two numbers, I mean two letters, reverse the letters. In this argument replace a with b and b with a . So then I get b minus the absolute value. The absolute value of b minus the absolute value of a is less than or equal to the absolute value of b minus a , which is the same as this.

And therefore-- let me multiply through by minus 1, and that tells me-- OK, so I have this is less than or equal to the absolute value of a minus b . I also have it's bigger than or equal to minus the absolute value of a minus b .

And therefore the absolute value of a minus the absolute value of b is less than or equal to absolute value of a minus b . So I it's taboo to write on the very back board, but I'm going to do it anyway. So that was the proof of the reverse triangle inequality. Here's the proof of the theorem before it.

It just follows from the reverse triangle inequality and that combination of those two theorems over there. We have that the absolute value of x sub n minus the absolute value of x . This is less than or equal to, by the reverse triangle inequality, the absolute value of x sub n minus x .

So this is by the reverse triangle inequality. And by assumption, this converges to 0 as n goes to infinity. So by the squeeze theorem, this goes to 0. And therefore the absolute value of x sub n converges to the absolute value of x . And that's it for that proof for this lecture and for this week.